



From Suspicion to Decision in Minutes

The world's first brain monitor for point-of-care seizure triage and treatment optimization


Seizures co-exist with many critical conditions and non-convulsive seizures are highly prevalent^{1,2}




25% OF ICU NEUROLOGICAL PATIENTS HAVE SEIZURES



Time to treatment is critical



Morbidity and mortality increase with prolonged NCSE^{3,4}



Response to first-line seizure treatment decreases over time⁵

Medical society guidelines recommend prompt EEG


NEUR CRITICAL

EEG should be initiated within **15-60 minutes** of suspected Status Epilepticus in all patients.⁶



We recommend **promptly performing and interpreting EEG** for the diagnosis of seizures in **all comatose patients after ROSC**.⁷





"Early access to EEG will lead to early detection, and hence, more effective treatment of seizures, which will in turn prevent refractory status epilepticus; neuronal injury; and potentially deleterious impacts on patient morbidity, mortality, and long-term outcome in terms of cognitive disability, overall neurologic function, and development of chronic epilepsy."

The DECIDE⁸ study is a multi-center prospective observational clinical study that evaluates the Clinical Impact of the Ceribell Rapid Response EEG.



Ceribell Point-of-Care EEG

Ceribell EEG Headband

Flexible headband accommodates various head types and head sizes



24/7 continuous bedside EEG monitoring and alerts



claritγ™

At the bedside, Clarity provides:

- First FDA cleared instantaneous bedside alert indicating suspected status epilepticus
- Continuous EEG monitoring and seizure burden display^{*9}

Remotely, Clarity provides:

- Prelabeled EEG making EEG reading more efficient
- First FDA cleared seizure burden trend for effective seizure management


*Seizure burden is defined by the American Clinical Neurophysiology Society (ACNS) as the percentage of time that EEG shows seizure activity.





Clinical Impact

- Changed clinical decisions in **40%** of cases.^{10,11}
- Spared **49%** of non-seizing patients from unnecessary medication.¹²
- Expedited ED disposition in 21% of patients.¹³



Enhancing your EEG capability to 24/7 on-site monitoring



- Minimize delays in treating non-convulsive status epilepticus
- Avoid unnecessary anti-seizure medications
- Avoid unnecessary patient transfers
- Reduce patient length of stay



Increased EEG access allows for accurate MS-DRG assignment for seizure-related CC/MCC¹⁹

Each EEG-triggered CC/MCC leads to \$5k to \$24K additional coding per case



Ceribell qualifies for the following EEG CPT codes

Routine EEG Codes - Codes include Professional and Technical components

Decording dynation	8+ Channels - No video		
Recording duration	Code	Work RVUs	
20-40 Minutes	95816*	1.08	
20-40 Minutes	95819*	1.08	
41-60 Minutes	95812	1.08	
61-119 Minutes	95813	1.63	

* 95816 Awake & Drowsy

95819 Awake & Asleep

Long-Term EEG Codes - PROFESSIONAL Component

Descuding duration	Referred to as	Time of report	8+ Channels - No video	
Recording duration			Code	Work RVUs
2-12 Hour	Partial day	Daily Report	95717	2.0
12-26 Hour	Full day	Daily Report	95719	3.0

Long-Term EEG Codes - TECHNICAL Component

De condicar donatica	Monitoring - 8+ Channels - No video			
Recording duration	None	Intermittent	Continuous	
2-12 Hour	95705	95706	95707	
12-26 Hour	95708	95709	95710	

Ceribell should be used for:

Complementing conventional EEG when tech/equipment is not available

Critical care EEG


- Emergency EEG in ED or ICU to detect status epilepticus
- · Prevention of treatment delays and of over-treatment


Ceribell should NOT be used for:

- Replacing long term video EEG monitoring
- Replacing conventional EEG to formally diagnose epilepsy



Ceribell's signal quality is equivalent to conventional EEG¹⁵






EEG using a circumferential 10-electrode montage meets the gold standard

An assessment of 169,510 EEGs showed that midline and parasagittal focal seizures were found in only 0.7% of EEGs.^{16,17}



Circumferential EEG had high diagnostic concordance (95%) with conventional EEG and had high agreement between EEG readers.¹⁸



Care Provider Benefits

24/7 rapid EEG access for faster and better critical care decisions.

Neurology EEG reading is covered using existing CPT codes.

Improve level of service and clinical care by reducing wait time for stat EEGs.

Improve quality of life for EEG technologists

Quickly respond to stat EEG requests and ability to triage to long term monitoring to best utilize your techs' time and equipment.

Improve quality of life for neurologists

Clarity continuous seizure monitoring can be used by non-neurologists during after-hours to avoid late calls to neurologists. Ceribell offers easy remote access to EEGs from any device with an internet connection. **Customize the workflow** to meet departmentspecific needs and patient selection criteria.

Grow neuro-service line without hiring additional EEG techs.

Increase neurology's profitability as Ceribell EEG is typically paid for by the operational budget of the department managing those patients (ICU, etc.), while neurology charges for reading fees.

CAUTION: FEDERAL (US) LAW RESTRICTS THIS DEVICE TO SALE BY OR ON THE ORDER OF A PHYSICIAN. REFER TO OPERATOR MANUAL AND LABELING FOR INDICATIONS, CONTRAINDICATIONS, WARNINGS, PRECAUTIONS AND INSTRUCTIONS FOR USE.

REFERENCES

 1. Herman, S.T., et al. (2015) J Clin Neurophysiol. 32(2):87-95

 2. Friedman, D., et al. (2009) Anesth Analg. 109(2):506-523

 3. Young, G.B., et al. (1996) Neurology. 47(1):83-89

 4. Beg, J.M., et al. (2017) J Med Econ. 20(1):45-53

 5. Lowenstein, D.H., et al. (1993) Neurology. 43(3 Pt 1): 483-488

 6. Brophy, G., et al. (2012) Neurocrit Care. 17(1):3-23

 7. Panchal, A.R., et al. (2020) Circulation. 142(suppl 2):S366-S468

 8. Vespa, P.M., et al. (2020) Crit Care Med. 48(9):1249-1257

 9. Hirsch, L.J., et al. (2013) J Clin Neurophysiol. 30(1):1-27

 10. Yazbeck, M., et al. (2019) J Neurosci Nurs. 51(6):308-312

 11. Hobbs, K., et al. (2018) Neurocrit Care. 29(2):302-312

 12. Kurup, D., et al. (2022) Epileptic Disorders. 24(5):1-7

 13. Wright, N., et al. (2021) EMJ, 38(12):923-926

 14. Kamousi, B., et al. (2022) American Epilepsy Society Abstract 1.475

 15. Kamousi, B., et al. (2019) Clin Neurophysiol Practice. 4:69-75

 16. Pedley, T.A., et al. (1981) Ann Neurol. 9:142-149

 17. Gururangan, K., et al. (2019) Neurocrit Care. 32(1):193-197

 18. Westover, M.B., et al. (2020) Neurocrit Care. 33(2):479-490

 19. Centers for Medicare & Medicaid Services - Medicare Inpatient Hospitals - by Geography and Service (2020) CMS.gov